arXiv:1005.3160v2 [math.GN] 20 May 2010

FUNNY PROBLEMS IN INTUITIVE TOPOLOGY

R. TAVAKOLI

Abstract. The goal of this article is to introduce some beautiful known riddles in intuitive topology; hoping to make at least some fun for the reader.

1. Funny problems in intuitive topology

The materials of this section are adapted from reference: [1].

Problem 1. ([1]) Show that the elastic body represented in figure 1 (a) can be deformed so as to become the one shown in 1 (b). In other words, were the human body elastic enough, after making linked rings with your index fingers and thumbs, you could move your hands apart without separating the joined fingertips.



Figure 1. Plot related to statement of problem 1 [1].

Problem 2. ([1]) A pretzel has two holes that "hold" a doughnut (see figure 2 (a)). Show that the pretzel can be deformed in such a way that one of its "handles" will unlink itself from the doughnut (figure 2 (b)).

Problem 3. ([1]) A circle is drawn on a pretzel with two holes (3 (a)). Show that it is possible to deform the pretzel so that the circle will be in the position represented in 3 (b).

Problem 4. ([1]) Show that a punctured tube from a bicycle tire can be turned inside out. More precisely, this would be possible if the rubber from which the tube is made were elastic enough. In real life it is impossible to turn a punctured tube inside out.

Problem 5. ([1]) Show that the fancy pretzel represented in 4 (a) can be deformed into the ordinary pretzel with two holes (4 (b)).

Department of Material Science and Engineering, Sharif University of Technology, Tehran, Iran, P.O. Box 11365-9466, tav@mehr.sharif.edu, rohtav@gmail.com.



FIGURE 2. Plot related to statement of problem 2 [1].



FIGURE 3. Plot related to statement of problem 3 [1].



FIGURE 4. Plot related to statement of problem 5 [1].

Do you can resolve these problems by yourself (please try and then go to the next page)?

Graphical solutions to above problems are introduced in figures 5 through 10.



FIGURE 5. graphical solution for problem 1 [1].



FIGURE 6. graphical solution for problem 2 [1].

Regarding to the problem 4, first we perform the deformations shown in figure 8. Then we can change the position of the obtained figure so that its "inside" (shown in white) becomes its "outside" (the shaded side of the surface) and vice-versa, simply by moving it as a rigid body in space until the hoop 1 occupies the position of the hoop 2. Once this is done, the previous deformations performed in reverse order result in the tube being turned inside out as required. Note that this

FIGURE 7. graphical solution for problem 3 [1].



FIGURE 8. graphical solution for problem 4 [1].

procedure interchanges the "parallel" and the "meridian" of the tube (see figure 9) [1].



FIGURE 9. graphical solution for problem 4 (continue) [1].

Regarding to the problem 5, first we perform the deformation shown in figure 10. The solid thus obtained (provided it is elastic) can clearly be def.ermed into the one shown in figure 1 (a). It now remains to apply the solution of Problem 1 (cf. figure 5)

Topologists call such objects isotopic (in space). More clearly, when we consider objects in space "up to deformations", i.e., we did not distinguish objects that can be deformed into each other by reshaping them, we called such objects as isotopic.

In the same way, the topology aid us to identify isotopic knots. For example all knots represented on the top row of figure **??** can be deformed into each other. Similarly, all knots represented on the bottom row of figure **??** can be deformed into each other. For solution readers are refereed to chapter 2 of [1].

REFERENCES

[1] VV Prasolov. *Intuitive topology.* Orient Blackswan.

FIGURE 10. graphical solution for problem 5 [1].

FIGURE 11. All knots represented on the top row are isotopic, and the same for the bottom row [1].

DEPARTMENT OF MATERIAL SCIENCE AND ENGINEERING, SHARIF UNIVERSITY OF TECHNOLOGY, TEHRAN, IRAN, P.O. BOX 11365-9466
E-mail address: tav@mehr.sharif.edu, rohtav@gmail.com